Home Health VNA IV Medication Clinical Fact Sheet							
IV Medication:	Infliximab / Remicade	Risk Level:					
Med Class:	Monoclonal Antibody	_					
	Common Uses:	Rheumatoid Arthritis, Crohn's Disease, Psoriatic Arthritis					
	Labs to Monitor:	CBC with Diff					
	Instructions/Precautions:	Hypersensitivity reaction; monitor BP and Pulse q 30min during infusion; CHF may occur or worsen; flu-like symptoms, serious infections and malignancies may occur; pre- meds frequently used; secondary infusion uses 1.2 filter; do not give if active infection					
	First Dose Allowed: Central Line Only: IV Push: Vesicant:						
	See Procedure Manual:	n/a					
Notes:	Only drugs listed as First Dose Allowed may be given in the home as a first dose and are						
	considered for a first dose on a case by case basis by the IV Program Manager						
	The IV Manager and/or Clinical Director must be c	onsulted before a first dose referral is accepted					
Risk Levels:	n/a = Routinely given; Clinician must be app 1= IV Program Mgr or Clinical Director ap 2= IV Program Mgr notification; Clinician	pproval before referral is accepted					

ľ

		Drug			
		Name			
		(generic/t		Common	
High	Risk	rade)	Class	Uses	Labs

1 Infliximab, Monoclona Rheumatoi CBC with D

Special Instructio ns	No 1st Dose	Central Line Only	IV Push		See Procedur e Manual
Hypersens	Х	0	0	0	0