

**Home Health VNA**  
**IV Medication Clinical Fact Sheet**

IV Medication: Hydration, PPN, TPN / Magnesium Risk Level: n/a  
Med Class: Hydration/ Nutrition

Common Uses: Short term hydration  
Labs to Monitor: CMP  
Instructions/Precautions: Flushing, sweating, hypotension, tetany

**\*\*Notes for all TPN Products: TPN patients need a double lumen for labs. If no double lumen available, draw peripheral labs; TPN patients must be on a stable cycle for 24 hours before discharge home; Diabetic patients must be on stable cycle for 48 hours before discharge home**

First Dose Allowed:	<table border="1"><tr><td>Y</td></tr></table>	Y
Y		
Central Line Only:	<table border="1"><tr><td>N</td></tr></table>	N
N		
IV Push:	<table border="1"><tr><td>N</td></tr></table>	N
N		
Vesicant:	<table border="1"><tr><td>N</td></tr></table>	N
N		

See Procedure Manual: n/a

Notes: Only drugs listed as First Dose Allowed may be given in the home as a first dose and are considered for a first dose on a case by case basis by the IV Program Manager  
The IV Manager and/or Clinical Director must be consulted before a first dose referral is accepted

Risk Levels: n/a = Routinely given; Clinician must be approved to administer IV medications  
1= IV Program Mgr or Clinical Director approval before referral is accepted  
2= IV Program Mgr notification; Clinicians must review Special Instructions