

Home Health VNA IV Medication Clinical Fact Sheet

IV Medication: Total Parental Nutrition(TPN) c or w/o Risk Level: 2
 Med Class: Nutrition

Common Uses: Short or long term nutrition support or
 Labs to Monitor: CBC c diff + Plts, CMP, CA, Mg, PO4, LFTs, PT
 Instructions/Precautions: Teach S/S hypo, hyperglycemia & dehydration.


First Dose Allowed:	N
Central Line Only:	Y
IV Push:	N
Vesicant:	N

See Procedure Manual: Inf Therapy TPN

****Notes for all TPN Products: TPN patients need a double lumen for labs. If no double lumen available, draw peripheral labs; TPN patients must be on a stable cycle for 24 hours before discharge home; Diabetic patients must be on stable cycle for 48 hours before discharge home**

Notes: Only drugs listed as First Dose Allowed may be given in the home as a first dose and are considered for a first dose on a case by case basis by the IV Program Manager
 The IV Manager and/or Clinical Director must be consulted before a first dose referral is accepted

Risk Levels: n/a = Routinely given; Clinician must be approved to administer IV medications
 1= IV Program Mgr or Clinical Director approval before referral is accepted
 2= IV Program Mgr notification; Clinicians must review Special Instructions



TPN
before