Home Health VNA IV Medication Clinical Fact Sheet

V Medication:	Eculizumab / Soliris	Risk Level: n/a
Med Class:	Monoclonal Antibody	_
	Common Uses: Labs to Monitor:	Paroxysmal Nocturnal Hemoglobinuia (PHN); Atypical Hemolytic Uremia Syndrome (aHUS) CBC, Chem 20, Urine Creatinine Ratio
	Instructions/Precautions:	For pediatric patients, dose is calculated by weight in kg; med is mixed in home; must be refrigerated and brought to room temp; withdraw required amount and transfer to IV bag; gently invert bag to mix; inspect for particulate and discoloration; infuse typically over 1 hour; hypersensitivity reactions have occurred; epinephrine and benedryl IM/IV are kept on hand
	First Dose Allowed: Central Line Only: IV Push: Vesicant:	N Y N N
	See Procedure Manual:	n/a
lotes:	Only drugs listed as First Dose Allowed may be given in the home as a first dose and are considered for a first dose on a case by case basis by the IV Program Manager The IV Manager and/or Clinical Director must be consulted before a first dose referral is accepted	
isk Levels:	n/a = Routinely given; Clinician must be approved to administer IV medications 1= IV Program Mgr or Clinical Director approval before referral is accepted 2= IV Program Mgr notification; Clinicians must review Special Instructions	