

| Criteria for Identification of Vulnerable Patients | Qualifier | Domain | Potential Contributing Team Members | | | | | | | | | | |
|------------------------------------------------------------------------------------------------------------------------|--------------|---------------|-------------------------------------|----|----|----|----|-----|-------|------|-------|----|---|
| | | | SN | PT | OT | ST | SW | HHA | Psych | WOCN | Palli | RD | |
| Repeated rehospitalizations (3 or more in 6 months) | Contributing | Clinical | x | x | x | | x | | | x | | x | |
| Uncontrolled or difficult to control pain | Contributing | Clinical | x | x | x | | | x | | | | x | |
| Inability to manage medications reliably with no willing or able caregiver | Contributing | Clinical | x | | x | | | | | x | | | |
| High Risk for Skin Breakdown (Braden score 12 or less) | Contributing | Clinical | x | x | x | | | x | | | x | | |
| Minimal VNA utilization (LUPA needs) but patient receives extensive supportive services from other providers or family | Primary | Clinical | x | x | x | | x | x | | | | | |
| Gradual or steady decline in clinical, functional and/or cognitive status | Contributing | Clinical | x | x | x | | | x | | | | x | |
| Weight loss / wasting syndrome | Contributing | Clinical | | | x | x | | x | | | | x | x |
| Multiple chronic conditions with deteriorating condition | Contributing | Clinical | x | x | x | | | | | | | x | |
| Chronic pressure ulcers | Contributing | Clinical | x | x | x | | | x | | | x | | x |
| Incontinence | Contributing | Clinical | x | x | x | | | x | | | x | | |
| Major functional limitation (i.e. bedbound, totally dependent for ADLs) | Primary | Functional | | x | x | | | x | | | | | |
| High Fall Risk (MAHC-10 score 4 or greater) or significant fall history | Contributing | Functional | x | x | x | | | x | | | x | | |
| Cognitive issues/ poor insight / self-neglect / abuse | Primary | Psycho-Social | x | | x | | x | x | x | | | | |
| Acute, severe psychiatric symptoms (delusions, delerium, major depression) | Contributing | Psycho-Social | | | x | | | x | x | | | | |
| Psychosocial issues / family dynamics impact negatively on plan of care or patient status | Primary | Psycho-Social | | | | | x | | x | | | x | |
| Suspected financial abuse | Contributing | Psycho-Social | | | | | x | | | | | | |
| Psychosocially isolated patient | Contributing | Psycho-Social | | | | | x | x | x | | | | |
| No Health Care Proxy, advanced directives, MOLST, DNR or unclear who is making decisions | Contributing | Psycho-Social | x | | | | x | | x | | | x | |
| Aggressive, intimidating caregiver | Contributing | Psycho-Social | x | | | | x | | x | | | | |
| Enrolled in PCA program or receiving 24-hour care from formal or informal caregiver that does not appear to meet needs | Contributing | Safety | x | | x | | x | | | | | | |
| Inadequate or absent support system (i.e. no primary caregiver, or unwilling/unable caregiver) | Contributing | Safety | x | | | | x | x | x | | | | |
| Unsafe environment (i.e. hoarding, aggressive animals, fire safety concern) | Primary | Safety | x | x | | | x | | x | | | | |
| Unable to leave home in an emergency | Primary | Safety | | x | | | x | x | | | | | |
| Criteria for filing protective services are present but ruled out or ineligible for services (age) | Primary | Safety | x | | | | x | | x | | | | |